

Universitext

UTX

Jean-Claude Hausmann

# Mod Two Homology and Cohomology

 Springer

---

# Universitext

## Series Editors

**Sheldon Axler, Vincenzo Capasso, Carles Casacuberta, Angus McIntyre, Kenneth Ribet, Claude Sabbah, Endre Süli and Wojbor A. Woźczynski**

*Universitext* is a series of textbooks that presents material from a wide variety of mathematical disciplines at master's level and beyond. The books, often well class-tested by their author, may have an informal, personal, even experimental approach to their subject matter. Some of the most successful and established books in the series have evolved through several editions, always following the evolution of teaching curricula, into very polished texts.

Thus as research topics trickle down into graduate-level teaching, first textbooks written for new, cutting-edge courses may make their way into *Universitext*.

More information about this series at <http://www.springer.com/series/223>

---

# Mod Two Homology and Cohomology



Springer

---

Jean-Claude Hausmann  
University of Geneva, Geneva, Switzerland

---

ISSN 0172-5939 e-ISSN 2191-6675

ISBN 978-3-319-09353-6 e-ISBN 978-3-319-09354-3  
DOI 10.1007/978-3-319-09354-3

Springer Cham Heidelberg New York Dordrecht London

Library of Congress Control Number: 2014944717

© Springer International Publishing Switzerland 2014

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission of information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media ([www.springer.com](http://www.springer.com))

---

# Contents

## 1 Introduction

## 2 Simplicial (Co)homology

### 2.1 Simplicial Complexes

### 2.2 Definitions of Simplicial (Co)homology

### 2.3 Kronecker Pairs

### 2.4 First Computations

#### 2.4.1 Reduction to Components

#### 2.4.2 $\mathbb{C}$ -Dimensional (Co)homology

#### 2.4.3 Pseudomanifolds

#### 2.4.4 Poincaré Series and Polynomials

#### 2.4.5 (Co)homology of a Cone

#### 2.4.6 The Euler Characteristic

#### 2.4.7 Surfaces

### 2.5 The Homomorphism Induced by a Simplicial Map

### 2.6 Exact Sequences

### 2.7 Relative (Co)homology

### 2.8 Mayer-Vietoris Sequences

### 2.9 Appendix A: An Acyclic Carrier Result

### 2.10 Appendix B: Ordered Simplicial (Co)homology

### 2.11 Exercises for Chapter 2

## 3 Singular and Cellular (Co)homologies

### 3.1 Singular (Co)homology

#### 3.1.1 Definitions

#### 3.1.2 Relative Singular (Co)homology

### **3.1.3 The Homotopy Property**

### **3.1.4 Excision**

### **3.1.5 Well Cofibrant Pairs**

### **3.1.6 Mayer-Vietoris Sequences**

### **3.2 Spheres, Disks, Degree**

### **3.3 Classical Applications of the $\text{mod } 2$ (Co)homology**

### **3.4 CW-Complexes**

### **3.5 Cellular (Co)homology**

### **3.6 Isomorphisms Between Simplicial and Singular (Co)homology**

### **3.7 CW-Approximations**

### **3.8 Eilenberg-MacLane Spaces**

### **3.9 Generalized Cohomology Theories**

### **3.10 Exercises for Chapter 3**

## **4 Products**

### **4.1 The Cup Product**

#### **4.1.1 The Cup Product in Simplicial Cohomology**

#### **4.1.2 The Cup Product in Singular Cohomology**

### **4.2 Examples**

#### **4.2.1 Disjoint Unions**

#### **4.2.2 Bouquets**

#### **4.2.3 Connected Sum(s) of Closed Topological Manifolds**

#### **4.2.4 Cohomology Algebras of Surfaces**

### **4.3 Two-Fold Coverings**

#### **4.3.1 $H^1$ , Fundamental Group and 2-Fold Coverings**

#### **4.3.2 The Characteristic Class**

### 4.3.3 The Transfer Exact Sequence of a $2$ -Fold Covering

### 4.3.4 The Cohomology Ring of $\mathbb{R}P^n$

## 4.4 Nilpotency, Lusterik-Schnirelmann Categories and Topological Complexity

## 4.5 The Cap Product

## 4.6 The Cross Product and the Künneth Theorem

## 4.7 Some Applications of the Künneth Theorem

### 4.7.1 Poincaré Series and Euler Characteristic of a Product

### 4.7.2 Slices

### 4.7.3 The Cohomology Ring of a Product of Spheres

### 4.7.4 Smash Products and Joins

### 4.7.5 The Theorem of Leray-Hirsch

### 4.7.6 The Thom Isomorphism

### 4.7.7 Bundles Over Spheres

### 4.7.8 The Face Space of a Simplicial Complex

### 4.7.9 Continuous Multiplications on $K(\mathbb{Z}_2, m)$

## 4.8 Exercises for Chapter 4

## 5 Poincaré Duality

### 5.1 Algebraic Topology and Manifolds

### 5.2 Poincaré Duality in Polyhedral Homology Manifolds

### 5.3 Other Forms of Poincaré Duality

#### 5.3.1 Relative Manifolds

#### 5.3.2 Manifolds with Boundary

#### 5.3.3 The Intersection Form

#### 5.3.4 Non Degeneracy of the Cup Product

#### 5.3.5 Alexander Duality

## **5.4 Poincaré Duality and Submanifolds**

---

### **5.4.1 The Poincaré Dual of a Submanifold**

### **5.4.2 The Gysin Homomorphism**

### **5.4.3 Intersections of Submanifolds**

### **5.4.4 The Linking Number**

### **5.5 Exercises for Chapter 5**

## **6 Projective Spaces**

### **6.1 The Cohomology Ring of Projective Spaces—Hopf Bundles**

### **6.2 Applications**

#### **6.2.1 The Borsuk-Ulam Theorem**

#### **6.2.2 Non-singular and Axial Maps**

### **6.3 The Hopf Invariant**

#### **6.3.1 Definition**

#### **6.3.2 The Hopf Invariant and Continuous Multiplications**

#### **6.3.3 Dimension Restrictions**

#### **6.3.4 Hopf Invariant and Linking Numbers**

### **6.4 Exercises for Chapter 6**

## **7 Equivariant Cohomology**

### **7.1 Spaces with Involution**

### **7.2 The General Case**

### **7.3 Localization Theorems and Smith Theory**

### **7.4 Equivariant Cross Products and Künneth Theorems**

### **7.5 Equivariant Bundles and Euler Classes**

### **7.6 Equivariant Morse-Bott Theory**

### **7.7 Exercises for Chapter 7**



## **8 Steenrod Squares**

### **8.1 Cohomology Operations**

### **8.2 Properties of Steenrod Squares**

### **8.3 Construction of Steenrod Squares**

### **8.4 Adem Relations**

### **8.5 The Steenrod Algebra**

### **8.6 Applications**

### **8.7 Exercises for Chapter 8**

## **9 Stiefel-Whitney Classes**

### **9.1 Trivializations and Structures on Vector Bundles**

### **9.2 The Class $w_1$ —Orientability**

### **9.3 The Class $w_2$ —Spin Structures**

### **9.4 Definition and Properties of Stiefel-Whitney Classes**

### **9.5 Real Flag Manifolds**

#### **9.5.1 Definitions and Morse Theory**

#### **9.5.2 Cohomology Rings**

#### **9.5.3 Schubert Cells and Stiefel-Whitney Classes**

### **9.6 Splitting Principles**

### **9.7 Complex Flag Manifolds**

### **9.8 The Wu Formula**

#### **9.8.1 Wu's Classes and Formula**

#### **9.8.2 Orientability and Spin Structures**

#### **9.8.3 Applications to 3-Manifolds**

#### **9.8.4 The Universal Class for Double Points**

### **9.9 Thom's Theorems**

## **9.9.1 Representing Homology Classes by Manifolds**

---

## **9.9.2 Cobordism and Stiefel-Whitney Numbers**

## **9.10 Exercises for Chapter 9**

## **10 Miscellaneous Applications and Developments**

## **10.1 Actions with Scattered or Discrete Fixed Point Sets**

## **10.2 Conjugation Spaces**

## **10.3 Chain and Polygon Spaces**

### **10.3.1 Definitions and Basic Properties**

### **10.3.2 Equivariant Cohomology**

### **10.3.3 Non-equivariant Cohomology**

### **10.3.4 The Inverse Problem**

### **10.3.5 Spatial Polygon Spaces and Conjugation Spaces**

## **10.4 Equivariant Characteristic Classes**

## **10.5 The Equivariant Cohomology of Certain Homogeneous Spaces**

## **10.6 The Kervaire Invariant**

## **10.7 Exercises for Chapter 10**

## **11 Hints and Answers for Some Exercises**

### **11.1 Exercises for Chapter 2**

### **11.2 Exercises for Chapter 3**

### **11.3 Exercises for Chapter 4**

### **11.4 Exercises for Chapter 5**

### **11.5 Exercises for Chapter 6**

### **11.6 Exercises for Chapter 7**

### **11.7 Exercises for Chapter 8**


### **11.8 Exercises for Chapter 9**

**References**

**Index**

---

# 1. Introduction

Jean-Claude Hausmann 

(1) University of Geneva, Geneva, Switzerland

 Jean-Claude Hausmann

Email: [jean-claude.hausmann@unige.ch](mailto:jean-claude.hausmann@unige.ch)

Mod 2 homology first occurred in 1908 in a paper of Tietze [196] (see also [40], pp. 41–42). Several results were first established using this  $\mathbb{Z}_2$  approach, like the linking number for submanifolds in  $\mathbb{R}^n$  (see Sect. 5.4.4), as well as Alexander duality [7]. One argument in favor of the choice of the  $\mathbb{Z}_2$  homology was its simplicity, as J.W. Alexander says in his introduction: “The theory of connectivity [homology] may be approached from two different angles depending on whether or not the notion of sense [orientation] is developed and taken into consideration. We have adopted the second and somewhat simpler point of view in this discussion in order to condense the necessary preliminaries as much as possible. A treatment involving the idea of sense would be somewhat more complicated but would follow along much the same lines.”

Besides being simpler than its integral counterpart,  $\mathbb{Z}_2$  homology sometimes gives new theorems. The first historical main example is the generalization of Poincaré duality to all closed manifolds, whether orientable or not, a result obtained by Veblen and Alexander in 1913 [200]. As a consequence, the Euler characteristic of a closed odd-dimensional manifold vanishes.

The discoveries of Stiefel-Whitney classes in 1936–1938 and of Steenrod squares in 1947–1950 gave  $\mathbb{Z}_2$  cohomology the status of a major tool in algebraic topology, providing for instance the theory of spin structures and Thom’s work on the cobordism ring.

These notes are an introduction, at graduate student’s level, of  $\mathbb{Z}_2$  (co)homology (there will be essentially no other). They include classical applications (Brouwer fixed point theorem, Poincaré duality, Borsuk-Ulam theorem, Smith theory, etc) and less classical ones (face spaces, topological complexity, equivariant Morse theory, etc). The cohomology of flag manifolds is treated in details, including for Grassmannians the relationship between Stiefel-Whitney classes and Schubert calculus. Some original applications are given in Chap. 10.

Our approach is different than that of classical textbooks, in which  $\mathbb{Z}_2$  (co)homology is just a particular case of (co)homology with arbitrary coefficients. Also, most authors start with a full account of homology before approaching cohomology. In these notes,  $\mathbb{Z}_2$  (co)homology is treated as a subject by itself and we start with cohomology and homology together from the beginning. The advantages of this approach are the following.

- The definition of a (co)chain is simple and intuitive: an (say, simplicial)  $m$ -cochain is a set of  $m$ -simplexes; an  $m$ -chain is a finite set of  $m$ -simplexes. The concept of cochain is simpler than that of chain (one less word in the definition...), more flexible and somehow more natural. We thus tend to consider cohomology as the main concept and homology as a (useful) tool for some

arguments.

- Working with  $\mathbb{Z}_2$  and its standard linear algebra is much simpler than working with  $\mathbb{Z}$ . For instance, the *Kronecker pairing* has an intuitive geometric interpretation occurring at the beginning which shows in an elementary way that cohomology is the dual of homology. Several computations, like the homology of surfaces, are quite easy and come early in the exposition. Also, the cohomology ring is *commutative*. The cup square  $\smile : H^k \rightarrow H^k \otimes H^k$  is a linear map and may be also non-trivial in odd degrees, leading to important invariants.
- The absence of sign and orientation considerations is an enormous technical simplification (even of importance in computer algorithms computing homology). With much lighter computations and technicalities, the ideas of proofs are more apparent.

We hope that these notes will be, for students and teachers, a complement or companion to textbooks like those of Hatcher [82] or Munkres [155]. From our teaching experience, starting with  $\mathbb{Z}_2$  (co)homology and taking advantage of its above mentioned simplicity is a great help to grasp the ideas of the subject. The technical difficulties of signs and orientations for finer theories, like integral (co)homology, may then be introduced afterwards, as an adaptation of the more intuitive  $\mathbb{Z}_2$  (co)homology.

*Not in this book* The following tools are not used in these notes.

- Augmented (co)chain complexes. The reduced cohomology  $\tilde{H}^*(X)$  is defined as  $\text{coker}(H^*(pt) \rightarrow H^*(X))$  for the unique map  $X \rightarrow pt$ .
- Simplicial approximation.
- Spectral sequences (except in the proof of Proposition 7.2.17).

Also, we do not use advanced homotopy tools, like spectra, completions, etc. Because of this, some prominent problems using  $\mathbb{Z}_2$  cohomology are only briefly surveyed, like the work by Adams on the Hopf-invariant-one problem (p. 353), the Sullivan's conjecture (pp. 240 and 353) and the Kervaire invariant (Sect. 10.6).


*Prerequisites* The reader is assumed to have some familiarity with the following subjects:

- general point set topology (compactness, connectedness, etc).
- elementary language of categories and functors.
- simple techniques of exact sequences, like the five lemma.
- elementary facts about fundamental groups, coverings and higher homotopy groups (not much used).
- elementary techniques of smooth manifolds.

## Acknowledgments

A special thank is due to Volker Puppe who provided several valuable suggestions and simplifications. Michel Zisman, Pierre de la Harpe, Samuel Tinguely and Matthias Franz have carefully read several sections of these notes. The author is also grateful for useful comments to Jim Davis, Rebecca Goldli, André Haefliger, Tara Holm, Allen Knutson, Jérôme Scherer, Dirk Schütz, Andras Szenes, Vladimir Turaev, Paul Turner, Claude Weber and Saïd Zarati.

## 2. Simplicial (Co)homology

Jean-Claude Hausmann 

(1) University of Geneva, Geneva, Switzerland

 **Jean-Claude Hausmann**

**Email:** [jean-claude.hausmann@unige.ch](mailto:jean-claude.hausmann@unige.ch)

Simplicial homology was invented by Poincaré in 1899 [162] and its  $\mathbb{Z}_2$  version, presented in this chapter, was introduced in 1908 by Tietze [196]. It is the simplest homology theory to understand and for finite complexes, it may be computed algorithmically. The  $\mathbb{Z}_2$  version permits rapid computations on easy but non-trivial examples, like spheres and surfaces (see Sect. 2.4).

Simplicial (co)homology is defined for a simplicial complex, but is an invariant of the homotopy type of its geometric realization (this result will be obtained in different ways using singular homology; see Sect. 3.6). The first section of this chapter introduces classical techniques of (abstract) simplicial complexes. Since simplicial homology was the only existing (co)homology theory until the 1930s, simplicial complexes played a predominant role in algebraic topology during the first third of the 20th century (see the Introduction of Sect. 5.1). Later developments of (co)homology theories, defined directly for topological spaces, made this combinatorial approach less crucial. However, simplicial complexes remain an efficient way to construct topological spaces, also largely used in computer science.

### 2.1 Simplicial Complexes

In this section we fix notations and recall some classical facts about (abstract) simplicial complexes. For more details, see [179, Chap. 3].

A *simplicial complex*  $K$  consists of

- a set  $V(K)$ , the set of *vertices* of  $K$ .
- a set  $\mathcal{S}(K)$  of finite non-empty subsets of  $V(K)$  which is closed under inclusion: if  $\sigma \in \mathcal{S}(K)$  and  $\tau \subset \sigma$ , then  $\tau \in \mathcal{S}(K)$ . We require that  $\{v\} \in \mathcal{S}(K)$  for all  $v \in V(K)$ .

An element  $\sigma$  of  $\mathcal{S}(K)$  is called a *simplex* of  $K$  (“simplexes” and “simplices” are admitted as plural of “simplex”; we shall use “simplexes”, in analogy with “complexes”). If  $\#\langle v \rangle = m + 1$ , we say that  $\sigma$  is of *dimension*  $m$  or that  $\sigma$  is an  $m$ -*simplex*. The set of  $m$ -simplexes of  $K$  is denoted by  $\mathcal{S}_m(K)$ . The set  $\mathcal{S}_0(K)$  of 0-simplexes is in bijection with  $V(K)$ , and we usually identify  $v \in V(K)$  with  $\langle v \rangle \in \mathcal{S}_0(K)$ . We say that  $K$  is of *dimension*  $< n$  if  $\mathcal{S}_m(K) = \emptyset$  for  $m > n$ , and that  $K$  is of *dimension*  $n$  if

(or  $m$ -dimensional) if it is of dimension  $< n$  but not of dimension  $< n - 1$ . A simplicial complex of dimension  $< 2$  is called a *simplicial graph*. A simplicial complex  $K$  is called *finite* if  $V(K)$  is a finite set.

If  $\sigma \in \mathcal{S}(K)$  and  $\tau \subset \sigma$ , we say that  $\tau$  is a *face* of  $\sigma$ . As  $\mathcal{S}(K)$  is closed under inclusion, it is determined by its subset  $\mathcal{S}_{\max}(K)$  of *maximal* simplices (if  $K$  is finite dimensional). A *subcomplex*  $L$  of  $K$  is a simplicial complex such that  $V(L) \subset V(K)$  and  $\mathcal{S}(L) \subset \mathcal{S}(K)$ . If  $S \subset \mathcal{S}(K)$  we denote by  $\bar{S}$  the *subcomplex generated by  $S$* , i.e. the smallest subcomplex of  $K$  such that  $S \subset \mathcal{S}(\bar{S})$ . The  $m$ -*skeleton*  $K^m$  of  $K$  is the subcomplex of  $K$  generated by the union of  $\mathcal{S}_k(K)$  for  $k \leq m$ .

Let  $\sigma \in \mathcal{S}(K)$ . We denote by  $\sigma$  the subcomplex of  $K$  formed by  $\sigma$  and all its faces ( $\{\bar{\sigma}\}$  in the above notation). The subcomplex  $\partial\sigma$  of  $\sigma$  generated by the proper faces of  $\sigma$  is called the *boundary* of  $\sigma$ .

### 2.1.1

*Geometric realization.* The *geometric realization*  $|K|$  of a simplicial complex  $K$  is, as a set, defined by

$$|K| := \{ \mu \in \mathcal{P}(V(K)) \mid \sum_{v \in V(K)} \mu(v) = 1 \text{ and } \mu^{-1}((0, 1]) \subset \mathcal{S}(K) \}$$

We can thus see  $|K|$  as the set of probability measures on  $V(K)$  which are supported by the simplices (this language is just used for comments and only in this section). There is a distance on  $|K|$  defined by

$$d(\mu, \nu) = \sqrt{\sum_{v \in V(K)} |\mu(v) - \nu(v)|^2}$$

which defines the *metric topology* on  $|K|$ . The set  $|K|$  with the metric topology is denoted by  $|K|_d$ . For instance, if  $\sigma \in \mathcal{S}_m(K)$ , then  $|\sigma|_d$  is isometric to the standard Euclidean simplex  $\Delta^m = \{(x_0, \dots, x_m) \in \mathbb{R}^{m+1} \mid x_i \geq 0 \text{ and } \sum x_i = 1\}$ .

However, a more used topology for  $|K|$  is the *weak topology*, for which  $A \subset |K|$  is closed if and only if  $A \cap |\sigma|_d$  is closed in  $|\sigma|_d$  for all  $\sigma \in \mathcal{S}(K)$ . The notation  $|K|_w$  stands for the set  $|K|$  endowed with the weak topology. A map  $f$  from  $|K|_w$  to a topological space  $X$  is then continuous if and only if its restriction to  $|\sigma|_d$  is continuous for each  $\sigma \in \mathcal{S}(K)$ . In particular, the identity  $|K| \rightarrow |K|_d$  is continuous which implies that  $|K|$  is Hausdorff. The weak and the metric topology coincide if and only if  $K$  is locally finite, that is each vertex is contained in a finite number of simplices. When  $K$  is not locally finite,  $|K|$  is not metrizable (see e.g. [179, Theorem 3.2.8]).

When a simplicial complex  $K$  is locally finite, has countably many vertices and is finite dimensional, it admits a *Euclidean realization*, i.e. an embedding of  $|K|_w$  into some Euclidean space  $\mathbb{R}^N$  which is piecewise affine. A map  $f: |K|_w \rightarrow \mathbb{R}^N$  is *piecewise affine* if, for each  $\sigma \in \mathcal{S}(K)$ , the restriction of  $f$  to  $|\sigma|_d$  is an affine map. Thus, for each simplex  $\sigma$ , the image of  $|\sigma|_d$  is an affine simplex.

of  $\mathbb{R}^n$ . If  $\dim K \leq n$ , such a realization exists in  $\mathbb{R}^{n+1}$  (see e.g. [179, Theorem 3.3.9]).

If  $v \in S(K)$  then  $|\bar{v}| \subset |K|$ . We call  $|\bar{v}|$  the *geometric simplex* associated to  $\sigma$ . Its *boundary* is  $|\partial v|$ . The space  $|\bar{v}| - |\partial v|$  is the *geometric open simplex* associated to  $\sigma$ . Observe that  $|K|$  is the disjoint union of its geometric open simplexes.

There is a natural injection  $\nu: V(K) \rightarrow |K|$  sending  $v$  to the Dirac measure with value 1 on  $v$ . We usually identify  $v$  with  $\nu(v)$ , seeing a simplex  $v$  as a point of  $|K|$  (a geometric vertex). In this way, a point  $\mu \in |K|$  may be expressed as a convex combination of (geometric) vertices:

$$\mu = \sum_{v \in V(K)} \mu(v)v. \quad (2.1)$$

### 2.1.2

Let  $K$  and  $L$  be simplicial complexes. Their *join* is the simplicial complex  $K * L$  defined by

$$(1) \quad V(K * L) = V(K) \cup V(L),$$

$$(2) \quad S(K * L) = S(K) \cup S(L) \cup \{\sigma \cup \tau \mid \sigma \in S(K) \text{ and } \tau \in S(L)\}.$$

Observe that, if  $\sigma \in S_r(K)$  and  $\tau \in S_s(L)$ , then  $\sigma \cup \tau \in S_{r+s+1}(K * L)$ . Also,  $\overline{\sigma \cup \tau} = \bar{\sigma} * \bar{\tau}$  and  $|K * L|$  the topological join of  $|K|$  and  $|L|$  (see p. 171).

### 2.1.3

*Stars, links, etc.* Let  $K$  be a simplicial complex and  $\sigma \in S(K)$ . The *star*  $St(\sigma)$  of  $\sigma$  is the subcomplex of  $K$  generated by all the simplexes containing  $\sigma$ . The *link*  $Lk(\sigma)$  of  $\sigma$  is the subcomplex of  $K$  formed by the simplexes  $\tau \in S(K)$  such that  $\tau \cap \sigma = \emptyset$  and  $\tau \cup \sigma \in S(K)$ . Thus,  $Lk(\sigma)$  is a subcomplex of  $St(\sigma)$  and

$$St(\sigma) = \bar{\sigma} * Lk(\sigma)$$

More generally, if  $L$  is a subcomplex of  $K$ , the *star*  $St(L)$  of  $L$  is the subcomplex of  $K$  generated by all the simplexes containing a simplex of  $L$ . The *link*  $Lk(L)$  of  $L$  is the subcomplex of  $K$  formed by the simplexes  $\tau \in S(St(L)) - S(L)$ . One has  $St(L) = L * Lk(L)$ . The *open star*  $Open(L)$  of  $L$  is the open neighbourhood of  $|L|$  in  $|K|$  defined by

$$Open(L) = \{\mu \in |K| \mid \mu(v) > 0 \text{ if } v \in V(L)\}$$

This is the interior of  $St(L)$  in  $|K|$ .

### 2.1.4



**Simplicial maps.** Let  $K$  and  $L$  be two simplicial complexes. A *simplicial map*  $f: K \rightarrow L$  is a map  $f: V(K) \rightarrow V(L)$  such that  $f(\sigma) \in \mathcal{S}(L)$  if  $\sigma \in \mathcal{S}(K)$ , i.e. the image of a simplex of  $K$  is a simplex of  $L$ .

Simplicial complexes and simplicial maps form a category, the *simplicial category*, denoted by  $\mathbf{Simp}$ .

A simplicial map  $f: K \rightarrow L$  induces a continuous map  $|f|: |K| \rightarrow |L|$  defined, for  $w \in V(L)$ , by

$$|f|(\mu)(w) = \sum_{v \in f^{-1}(w)} \mu(v)$$

In other words,  $|f|(\mu)$  is the pushforward of the probability measure  $\mu$  on  $|K|$ . The geometric realization is thus a covariant functor from the simplicial category  $\mathbf{Simp}$  to the topological category  $\mathbf{Top}$  of topological spaces and continuous maps.

### 2.1.5

**Components.** Let  $K$  be a simplicial complex. We define an equivalence relation on  $V(K)$  by saying that  $v \sim v'$  if there exists  $x_0, \dots, x_m \in V(K)$  with  $x_0 = v$ ,  $x_m = v'$  and  $\{x_i, x_{i+1}\} \in \mathcal{S}(K)$ . A maximal subcomplex  $L$  of  $K$  such that  $V(L)$  is an equivalence class is called a *component* of  $K$ . The set of components of  $K$  is denoted by  $\pi_0(K)$ . As the vertices of a simplex are all equivalent,  $K$  is the disjoint union of its components and  $\pi_0(K)$  is in bijection with  $V(K)/\sim$ . The relationship with  $\pi_0(|K|)$  the set of (path)-components of the topological space  $|K|$ , is the following.

#### Lemma 2.1.6

The natural injection  $j: V(K) \rightarrow |K|$  descends to a bijection  $\bar{j}: \pi_0(K) \xrightarrow{\cong} \pi_0(|K|)$ .

#### Proof

The definition of the relation  $\sim$  makes clear that  $j$  descends to a map  $\bar{j}: \pi_0(K) \rightarrow \pi_0(|K|)$ . Any point of  $|K|$  is joinable by a continuous path to some vertex  $j(v)$ . Hence,  $\bar{j}$  is surjective. To check the injectivity of  $\bar{j}$ , let  $v, v' \in V(K)$  with  $\bar{j}(v) = \bar{j}(v')$ . There exists then a continuous path  $c: [0, 1] \rightarrow |K|$  with  $c(0) = j(v)$  and  $c(1) = j(v')$ . Consider the open cover  $\{\text{Ost}(w) \mid w \in V(K)\}$  of  $|K|$ . By compactness of  $[0, 1]$ , there exists  $n \in \mathbb{N}$  and vertices  $v_0, \dots, v_{n-1} \in V(K)$  such that  $c([k/n, (k+1)/n]) \subset \text{Ost}(v_k)$  for all  $k = 0, \dots, n-1$ . As  $c(0) = j(v)$  and  $c(1) = j(v')$ , one deduces that  $v_0 = v$  and  $v_{n-1} = v'$ . For  $0 < k \leq n-1$  one has  $c([k/n, (k+1)/n]) \subset \text{Ost}(v_{k-1}) \cap \text{Ost}(v_k)$ . This implies that  $\{v_{k-1}, v_k\} \in \mathcal{S}(K)$  for all  $k = 1, \dots, n-1$ , proving that  $v \sim v'$ .  $\square$

A simplicial complex is called *connected* if it is either empty or has one component. Note that  $K$  is locally path-connected for any simplicial complex  $K$ . Indeed, any point has a neighborhood of the form  $|\text{St}(v)|$  for some vertex  $v$ , and  $|\text{St}(v)|$  path-connected. Therefore,  $K$  is path-connected if and only if  $|K|$  is connected. Using Lemma 2.1.6, this proves the following lemma.

#### Lemma 2.1.7

Let  $K$  be a simplicial complex. Then  $K$  is connected if and only if  $|K|$  is a connected space.

Finally, we note the functoriality of  $\pi_0$ . Let  $f: K \rightarrow L$  be a simplicial map. If  $v \sim v'$  for  $v, v' \in V(K)$ , then  $f(v) \sim f(v')$ , so  $f$  descends to a map  $\pi_0 f: \pi_0(K) \rightarrow \pi_0(L)$ . If  $f: K \rightarrow L$  and  $g: L \rightarrow M$  are two simplicial maps, then  $\pi_0(g \circ f) = \pi_0 g \circ \pi_0 f$ . Also,  $\pi_0 \text{id}_K = \text{id}_{\pi_0(K)}$ . Thus,  $\pi_0$  is a covariant functor from the simplicial category **Simp** to the category **Set** of sets and maps.

### 2.1.8

**Simplicial order.** A simplicial order on a simplicial complex  $L$  is a partial order  $\leq$  on  $V(L)$  such that each simplex is totally ordered. For example, a total order on  $V(L)$ , as in examples where vertices are labeled by integers, is a simplicial order. A simplicial order always exists, as a consequence of the well-ordering theorem.

### 2.1.9

**Triangulations.** A triangulation of a topological space  $X$  is a homeomorphism  $h: |K| \rightarrow X$ , where  $K$  is a simplicial complex. A topological space is triangulable if it admits a triangulation. It will be useful to have a good process to triangulate some subspaces of  $\mathbb{R}^n$ . A compact subspace  $A$  of  $\mathbb{R}^n$  is a convex cell if it is the set of solutions of families of affine equations and inequalities

$$f_i(x) = 0, i = 1, \dots, r \quad \text{and} \quad g_j(x) \geq 0, j = 1, \dots, s.$$

A face  $B$  of  $A$  is a convex cell obtained by replacing some of the inequalities  $g_j \geq 0$  by the equations  $g_j = 0$ . The dimension of  $B$  is the dimension of the smallest affine subspace of  $\mathbb{R}^n$  containing  $B$ . A vertex of  $A$  is a cell of dimension 0. By induction on the dimension, one proves that a convex cell is the convex hull of its vertices (see e.g. [138, Theorem 5.2.2]).

A convex-cell complex  $P$  is a finite union of convex cells in  $\mathbb{R}^n$  such that:

- (i) if  $A$  is a cell of  $P$ , so are the faces of  $A$ ;
- (ii) the intersection of two cells of  $P$  is a common face of each of them.

The dimension of  $P$  is the maximal dimension of a cell of  $P$ . The  $r$ -skeleton  $P^r$  is the subcomplex formed by the cells of dimension  $\leq r$ . The 0-skeleton coincides with the set  $V(P)$  of vertices of  $P$ .

A partial order  $\leq$  on  $V(P)$  is an affine order for  $P$  if any subset  $R \in V(P)$  formed by affinely independent points is totally ordered. For instance, a total order on  $V(P)$  is an affine order. The following lemma is a variant of [104, Lemma 1.4].

#### Lemma 2.1.10

Let  $P$  be a convex-cell complex. An affine order  $\leq$  for  $P$  determines a triangulation  $h_{\leq}: |L_{\leq}| \xrightarrow{\cong} P$ ,

where  $L_\infty$  is a simplicial complex with  $V(L_\infty) = V(P)$ . The homeomorphism  $h_\infty$  is piecewise affine and  $\leq$  is a simplicial order on  $L_\infty$ .

### Proof

The order  $\leq$  being chosen, we drop it from the notations. For each subcomplex  $Q$  of  $P$ , we shall construct a simplicial complex  $L(Q)$  and a piecewise affine homeomorphism  $h_Q: |L(Q)| \rightarrow Q$  such that,

(i)  $V(L(Q)) = V(Q)$ ;

(ii) if  $Q' \subset Q$ , then  $L(Q') \subset L(Q)$  and  $h_{Q'}$  is the restriction of  $h_Q$  to  $L(Q')$ .

The case  $Q = P$  will prove the lemma. The construction is by induction on the dimension of  $Q$ , setting  $L(Q) = Q$  and  $h_Q = \text{id}$  if  $\dim Q = 0$ .

Suppose that  $L(Q)$  and  $h_Q$  have been constructed, satisfying (i) and (ii) above, for each subcomplex  $Q$  of  $P$  of dimension  $< k - 1$ . Let  $A$  be a  $k$ -cell of  $K$  with minimal vertex  $a$ . Then  $A$  is the topological cone, with cone-vertex  $a$ , of the union  $B$  of faces of  $A$  not containing  $a$ . The triangulation  $h_B: |L(B)| \rightarrow |B|$  being constructed by induction hypothesis, define  $L(A)$  to be the join  $L(B) * \{a\}$  and  $h_A$  to be the unique piecewise affine extension of  $h_B$ . Observe that, if  $C$  is a face of  $A$ , then  $h_C$  is the restriction to  $L(C)$  of  $h_A$ . Therefore, this process may be used for each  $k$ -cell of  $P$  to construct  $h_Q: |L(Q)| \rightarrow Q$  for each subcomplex  $Q$  of  $P$  with  $\dim Q \leq k$ .  $\square$

### 2.1.11

**Subdivisions.** Let  $Z$  be a set and  $\mathcal{A}$  be a family of subsets of  $Z$ . A simplicial complex  $L$  such that

(a)  $V(L) \subset Z$ ;

(b) for each  $\sigma \in S(L)$  there exists  $A \in \mathcal{A}$  such that  $\sigma \subset A$ ;

is called a  $(Z, \mathcal{A})$ -simplicial complex, or a  $Z$ -simplicial complex supported by  $\mathcal{A}$ .

Let  $K$  be a simplicial complex. Let  $N$  be a  $(|K|, \mathcal{GS}(K))$ -simplicial complex, where

$$\mathcal{GS}(K) = \{|\sigma| \mid \sigma \in S(K)\}$$

is the family of geometric simplexes of  $K$ . A continuous map  $i: |N| \rightarrow |K|$  is associated to  $N$ , defined

by

$$j(\mu) = \sum_{\omega \in V(\mu)} \mu(\omega)\omega$$

In other words,  $j$  is the piecewise affine map sending each vertex of  $N$  to the corresponding point in  $|K|$ . A subdivision of a simplicial complex  $K$  is a  $(|K|, \mathcal{G}\mathcal{S}(K))$ -simplicial complex  $N$  for which the associated map  $j: |N| \rightarrow |K|$  is a homeomorphism (in other words,  $j$  is a triangulation of  $|K|$ ).

Let  $N$  be a  $(|K|, \mathcal{G}\mathcal{S}(K))$ -simplicial complex for a simplicial complex  $K$ . If  $L$  is a subcomplex of  $K$ , then

$$N_L = \{\sigma \in \mathcal{S}(N) \mid \sigma \subset L\}$$

is a  $(|L|, \mathcal{G}\mathcal{S}(L))$ -simplicial complex. Its associated map  $j_L: |N_L| \rightarrow |L|$  is the restriction of  $j$  to  $|L|$ .

The following Lemma is useful to recognize a subdivision (compare [179, Chap. 3, Sect. 3, Theorem 4]).

### Lemma 2.1.12

Let  $N$  be a  $(|K|, \mathcal{G}\mathcal{S}(K))$ -simplicial complex. Then  $N$  is a subdivision of  $K$  if and only if, for each  $\tau \in \mathcal{S}(K)$ , the simplicial complex  $N_\tau$  is finite and  $j_\tau: |N_\tau| \rightarrow |\tau|$  is bijective.

#### Proof

If  $N$  is a subdivision of  $K$ , then  $j_\tau$  is bijective since  $j$  is a homeomorphism. Also,  $|N_\tau| = j^{-1}(|\tau|)$  is compact, so  $N_\tau$  is finite.

Conversely, the fact that  $j_\tau$  is bijective for each  $\tau \in \mathcal{S}(K)$  implies that the continuous map  $j$  is bijective. If  $N_\tau$  is finite, then  $j_\tau$  is a continuous bijection between compact spaces, hence a homeomorphism. This implies that the map  $j^{-1}$ , restricted to each geometric simplex, is continuous. Therefore,  $j^{-1}$  is continuous since  $K$  is endowed with the weak topology.  $\square$

Seeing  $V(K)$  as a subset of  $|K|$ , we get the following corollary.

### Corollary 2.1.13

Let  $N$  be a subdivision of  $K$ . Then  $V(K) \subset V(N)$ .

A useful systematic subdivision process is the barycentric subdivision. Let  $\sigma \in \mathcal{S}_m(K)$  be an  $m$ -simplex of a simplicial complex  $K$ . The *barycenter*  $\hat{\sigma} \in |K|$  of  $\sigma$  is defined by

$$\hat{\sigma} = \frac{1}{m+1} \sum_{\alpha \in \mathbb{S}_m} v_\alpha.$$

The *barycentric subdivision*  $K'$  of  $K$  is the  $(|K|, \mathcal{G}\mathcal{S}(K))$ -simplicial complex where

- $V(K') = \{\hat{\sigma} \in |K| \mid \sigma \in \mathcal{S}(K)\}$ ;
- $\{\hat{\sigma}_0, \dots, \hat{\sigma}_m\} \in \mathcal{S}_m(K')$  whenever  $\sigma_0 \subset \dots \subset \sigma_m$  ( $\sigma_i \neq \sigma_j$  if  $i \neq j$ ).

Using Lemma 2.1.12, the reader can check that  $K'$  is a subdivision of  $K$ . Observe that the partial order " $\leq$ " defined by

$$\sigma \leq \tau \iff \sigma \subset \tau \tag{2.1}$$

is a simplicial order on  $K'$ .

## 2.2 Definitions of Simplicial (Co)homology

Let  $K$  be a simplicial complex. In this section, we give the definitions of the homology  $H_*(K)$  and cohomology  $H^*(K)$  of  $K$  under the various and peculiar forms available when the coefficients are in the field  $\mathbb{Z}_2 = \{0, 1\}$ .

### Definition 2.2.1

(subset definitions)

- (a) An  $m$ -cochain is a subset of  $\mathcal{S}_m(K)$ .
- (b) An  $m$ -chain is a finite subset of  $\mathcal{S}_m(K)$ .

The set of  $m$ -cochains of  $K$  is denoted by  $C^m(K)$  and that of  $m$ -chains by  $C_m(K)$ . By identifying  $\sigma \in \mathcal{S}_m(K)$  with the singleton  $\{\sigma\}$ , we see  $\mathcal{S}_m(K)$  as a subset of both  $C_m(K)$  and  $C^m(K)$ . Each subset of  $\mathcal{S}_m(K)$  is determined by its characteristic function  $\chi_A: \mathcal{S}_m(K) \rightarrow \mathbb{Z}_2$ , defined by

$$\chi_A(\sigma) = \begin{cases} 1 & \text{if } \sigma \in A \\ 0 & \text{otherwise} \end{cases}$$

This gives a bijection between subsets of  $\mathcal{S}_m(K)$  and functions from  $\mathcal{S}_m(K)$  to  $\mathbb{Z}_2$ . We see such a function as a colouring (0 = white and 1 = black). The following "colouring definition" is equivalent to the subset definition:

### Definition 2.2.2

(colouring definitions)

- (a) An  $m$ -cochain is a function  $\alpha: \mathcal{S}_m(K) \rightarrow \mathbb{Z}_2$ .
- (b) An  $m$ -chain is a function  $\alpha: \mathcal{S}_m(K) \rightarrow \mathbb{Z}_2$  with finite support.

The colouring definition is used in low-dimensional graphical examples to draw (co)chains in black (bold lines for 1-(co)chains).

Definition 2.2.2 endow  $C^m(K)$  and  $C_m(K)$  with a structure of a  $\mathbb{Z}_2$ -vector space. The singletons provide a basis of  $C_m(K)$ , in bijection with  $\mathcal{S}_m(K)$ . Thus, Definition 2.2.2b is equivalent to

### Definition 2.2.3

$C_m(K)$  is the  $\mathbb{Z}_2$ -vector space with basis  $\mathcal{S}_m(K)$ :

$$C_m(K) = \bigoplus_{\sigma \in \mathcal{S}_m(K)} \mathbb{Z}_2 \sigma$$

We shall pass from one of Definitions 2.2.1, 2.2.2 or 2.2.3 to another without notice; the context

usually prevents ambiguity. We consider  $C_*(K) = \bigoplus_{m \in \mathbb{N}} C_m(K)$  and  $C^*(K) = \bigoplus_{m \in \mathbb{N}} C^m(K)$  as graded  $\mathbb{Z}_2$ -vector spaces. The convention  $C_{-1}(K) = C^{-1}(K) = 0$  is useful.

We now define the *Kronecker pairing* on (co)chains

$$C^m(K) \times C_m(K) \xrightarrow{\langle \cdot, \cdot \rangle} \mathbb{Z}_2$$

by the equivalent formulae

$$\begin{aligned} \langle a, \alpha \rangle &= \#(a \cap \alpha) \pmod{2} && \text{using Definition 2.2.1a and b} \\ &= \sum_{\sigma \in a} \alpha(\sigma) && \text{using Definitions 2.2.1a and 2.2.2b} \\ &= \sum_{\sigma \in S_m(K)} \alpha(\sigma) \alpha(\sigma) && \text{using Definitions 2.2.2a and b.} \end{aligned} \tag{2.2.1}$$

#### Lemma 2.2.4

The Kronecker pairing is bilinear and the map  $\alpha \mapsto \langle a, \cdot \rangle$  is an isomorphism between  $C^m(K)$  and  $C_m(K)^\# = \text{hom}(C_m(K), \mathbb{Z}_2)$ .

#### Proof

The bilinearity is obvious from the third line of Eq. (2.2.1). Let  $0 \neq a \in C^m(K)$ . This means that, as a subset of  $S_m(K)$ ,  $a$  is not empty. If  $\sigma \subset a$ , then  $\langle a, \sigma \rangle \neq 0$ , which proves the injectivity of  $\alpha \mapsto \langle a, \cdot \rangle$ . As for its surjectivity, let  $h \in \text{hom}(C_m(K), \mathbb{Z}_2)$ . Using the inclusion  $S_m(K) \hookrightarrow C_m(K)$  given by  $\tau \mapsto \{\tau\}$ , define

$$a = \{\tau \in S_m(K) \mid h(\tau) = 1\}.$$

For each  $\sigma \in S_m(K)$  the equation  $h(\sigma) = \langle a, \sigma \rangle$  holds true. As  $S_m(K)$  is a basis of  $C_m(K)$ , this implies that  $h = \langle a, \cdot \rangle$ .  $\square$

We now define the boundary and coboundary operators. The *boundary operator*  $\partial: C_m(K) \rightarrow C_{m-1}(K)$  is the  $\mathbb{Z}_2$ -linear map defined by

$$\partial(\sigma) = \{(m-1) \text{ faces of } \sigma\} = S_{m-1}(\sigma), \sigma \in S_m(K). \tag{2.2.2}$$

Formula (2.2.2) is written in the language of Definition 2.2.1b. Using Definition 2.2.3, we get

$$\partial(v) = \sum_{\tau \in S_{m-1}(\bar{v})} \tau. \tag{2.2.3}$$

The *coboundary operator*  $\delta: C^m(K) \rightarrow C^{m+1}(K)$  is defined by the equation

$$\langle \delta a, \alpha \rangle = \langle a, \partial \alpha \rangle. \tag{2.2.4}$$

The last equation indeed defines  $\delta$  by Lemma 2.2.4 and  $\delta$  may be seen as the Kronecker adjoint of  $\partial$ . In particular, if  $\sigma \in S_m(K)$  and  $\tau \in S_{m-1}(K)$  then

$$\tau \in \partial(\sigma) \Leftrightarrow \tau \subset \sigma \Leftrightarrow \sigma \in \delta(\tau). \tag{2.2.5}$$

The first equivalence determines the operator  $\delta$  since  $S_m(K)$  is a basis for  $C_m(K)$ . The second equivalence determines  $\delta$  if  $S_{m-1}(K)$  is finite. Note that the definition of  $\delta$  may also be given as follows; if  $a \in C^m(K)$ , then

$\delta(\alpha) = \{\sigma \in S_{m+1}(K) \mid \#(\alpha \cap \delta(\sigma)) \text{ is odd}\}$   
 i.e.  $\sigma \in S_m(K)$ . Each  $\tau \in S_{m-1}(K)$  with  $\tau \subset \sigma$  belongs to the boundary of exactly two  $(m-1)$ -simplexes of  $\sigma$ . Using Eq. (2.2.3), this implies that  $\partial \cdot \partial = 0$ . By Eq. (2.2.4) and Lemma 2.2.4, we get  $\delta \circ \delta = 0$ . We define the  $\mathbb{Z}_2$ -vector spaces

- $Z_m(K) = \ker(\delta : C_m(K) \rightarrow C_{m-1}(K))$ , the  $m$ -cycles of  $K$ .
- $B_m(K) = \text{image}(\partial : C_{m+1}(K) \rightarrow C_m(K))$ , the  $m$ -boundaries of  $K$ .
- $Z^m(K) = \ker(\delta : C^m(K) \rightarrow C^{m-1}(K))$ , the  $m$ -cocycles of  $K$ .
- $B^m(K) = \text{image}(\delta : C^{m-1}(K) \rightarrow C^m(K))$ , the  $m$ -coboundaries of  $K$ .

For example, Fig. 2.1 shows a triangulation  $K$  of the plane, with  $V(K) = \mathbb{Z} \times \mathbb{Z}$ . The bold line is a cocycle  $\alpha$  which is a coboundary:  $\alpha = \delta\beta$ , with  $\beta = \{\{(m, n)\} \mid (m, n) \in V(K) \text{ and } m < 0\}$ , drawn in bold dots.




Fig. 2.1 A triangulation  $K$  of the plane, with  $V(K) = \mathbb{Z} \times \mathbb{Z}$

Since  $\partial \circ \delta = 0$  and  $\delta \circ \delta = 0$ , one has  $B_m(K) \subset Z_m(K)$  and  $B^m(K) \subset Z^m(K)$ . We form the quotient vector spaces

- $H_m(K) = Z_m(K)/B_m(K)$ , the  $m$ th-homology vector space of  $K$ .
- $H^m(K) = Z^m(K)/B^m(K)$ , the  $m$ th-cohomology vector space of  $K$ .

As for the (co)chains, the notations  $H_*(K) = \bigoplus_{m \in \mathbb{N}} H_m(K)$  and  $H^*(K) = \bigoplus_{m \in \mathbb{N}} H^m(K)$  stand for the (co)homology seen as graded  $\mathbb{Z}_2$ -vector spaces. By convention,  $H_{-1}(K) = H^{-1}(K) = 0$ . Also, the homology and the cohomology are in duality via the Kronecker pairing:

### Proposition 2.2.5

(Kronecker duality) The Kronecker pairing on (co)chains induces a bilinear map

$$H^m(K) \times H_m(K) \xrightarrow{\langle \cdot, \cdot \rangle} \mathbb{Z}_2$$

Moreover, the correspondence  $a \mapsto \langle a, \cdot \rangle$  is an isomorphism

$$H^m(K) \xrightarrow{\mathbf{k}} \text{Hom}(H_m(K), \mathbb{Z}_2).$$

### Proof

Instead of giving a direct proof, which the reader may do as an exercise, we will take advantage of the more general setting of *Kronecker pairs*, developed in the next section. In this way, Proposition 2.2.5 follows from Proposition 2.3.5.  $\square$

## 2.3 Kronecker Pairs

All the vector spaces in this section are over an arbitrary fixed field  $\mathbb{F}$ . The dual of a vector space  $V$  is denoted by  $V^\#$ .

A *chain complex* is a pair  $(C_*, \partial)$ , where

- $C_*$  is a graded vector space  $C_* = \bigoplus_{m \in \mathbb{N}} C_m$ . We add the convention that  $C_{-1} = 0$ .
- $\partial : C_* \rightarrow C_*$  is a linear map of degree  $-1$ , i.e.  $\partial(C_m) \subset C_{m-1}$ , satisfying  $\partial \circ \partial = 0$ . The operator  $\partial$  is called the *boundary* of the chain complex.

A *cochain complex* is a pair  $(C^*, \delta)$ , where

- $C^*$  is a graded vector space  $C^* = \bigoplus_{m \in \mathbb{N}} C^m$ . We add the convention that  $C^{-1} = 0$ .
- $\delta : C^* \rightarrow C^*$  is a linear map of degree  $+1$ , i.e.  $\delta(C^m) \subset C^{m+1}$ , satisfying  $\delta \circ \delta = 0$ . The operator  $\delta$  is called the *coboundary* of the cochain complex.

A *Kronecker pair* consists of three items:

(a) a chain complex  $(C_*, \partial)$ .

(b) a cochain complex  $(C^*, \delta)$ .

(c) a bilinear map

$$C^m \times C_m \xrightarrow{\langle \cdot, \cdot \rangle} \mathbb{F}$$

satisfying the equation

$$\langle \delta a, \alpha \rangle = \langle a, \delta \alpha \rangle, \tag{2.3}$$

for all  $a \in C^m$  and  $\alpha \in C_{m+1}$  and all  $m \in \mathbb{N}$ . Moreover, we require that the map  $\mathbf{k} : C^m \rightarrow C_m^\#$ , given by  $\mathbf{k}(a) = \langle a, \cdot \rangle$ , is an isomorphism.

### Example 2.3.1

Let  $K$  be a simplicial complex. Its simplicial (co)chain complexes  $(C^*(K), \delta)$ ,  $(C_*(K), \partial)$ , together with



- [read online From Lowbrow to Nobrow pdf, azw \(kindle\), epub](#)
- [read The Wire, Issue 380 \(October 2015\) here](#)
- [click \*\*Das Beste an HTML & CSS pdf, azw \(kindle\)\*\*](#)
- [read online The J Curve: A New Way to Understand Why Nations Rise and Fall](#)
  
- <http://aneventshop.com/ebooks/From-Lowbrow-to-Nobrow.pdf>
- <http://www.mmastyles.com/books/The-Wire--Issue-380--October-2015-.pdf>
- <http://interactmg.com/ebooks/The-Road-to-Bedlam--Courts-of-the-Feyre--Book-2-.pdf>
- <http://interactmg.com/ebooks/A-Companion-to-Latin-American-Literature-and-Culture.pdf>